


(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 January 2001 (18.01.2001)

PCT


(10) International Publication Number
WO 01/04100 A1

- (51) International Patent Classification⁷: C07D 239/42, C07C 229/34, 311/05
- (74) Common Representative: LONZA AG; Münchensteinerstrasse 38, CH-4052 Basel (CH).
- (21) International Application Number: PCT/EP00/06099
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 30 June 2000 (30.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|-------------------------------|----|
| 99113711.8 | 13 July 1999 (13.07.1999) | EP |
| 99120417.3 | 14 October 1999 (14.10.1999) | EP |
| 60/185,465 | 28 February 2000 (28.02.2000) | US |
| 60/185,371 | 28 February 2000 (28.02.2000) | US |
| 00106303.1 | 23 March 2000 (23.03.2000) | EP |
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): LONZA AG [CH/CH]; Münchensteinerstrasse 38, CH-4052 Basel (CH).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): VEITH, Ulrich [DE/CH]; Weingartenweg 16, CH-3930 Visp (CH).
- Published:**
- With international search report.
 - Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*


(54) Title: PROCESS FOR PREPARING 2-AMINO-4-(4-FLUOROPHENYL)-6-ALKYLPYRIMIDINE-5-CARBOXYLATE



(I)



(II)



(III)




(VI)

(57) Abstract: Process for the preparation of compounds of general formula (I), in which R is hydrogen or a group of formula -SO₂R¹, R¹ is C₁₋₆-alkyl; R² is hydrogen or C₁₋₆-alkyl; R³ is C₁₋₆-alkyl; R⁴ is C₁₋₆-alkyl, characterized in that, in a first stage, a compound of formula (II), in which R³ and R⁴ have the abovementioned meaning, is reacted in the presence of a Lewis acid with 4-fluorobenzonitrile to give a compound of general formula (III), in which R³ and R⁴ have the abovementioned meaning and in a second stage the compound of formula (III) obtained is reacted with the compound of formula (IV), in which R and R² have the abovementioned meaning, to give the final product of formula (I).

PROCESS FOR PREPARING 2-AMINO-4-(4-FLUOROPHENYL)-6-
ALKYLPYRIMIDINE-5-CARBOXYLATE

Description:

The present invention relates to a novel
process for the preparation of compounds of the general
5 formula



in which R, R², R³ and R⁴ have the meaning below.

Compounds of the formula I are important
10 intermediates for the preparation of pharmaceutically
active compounds, for example of HMG-Co A reductase
inhibitors. Japanese Patent publication JP-A 06 256318,
and Watanabe M. et al., *Bioorg. Med. Chem.* 1997,
Vol. 5, No. 2, 437-444 describe processes for the
15 preparation of compounds of the formula I.

The process described in JP-A 06 256318 has the
disadvantage that three stages are needed in order to
prepare 2-amino-4-(4-fluorophenyl)-6-isopropyl-pyri-
20 midine-5-carboxylic acid.

A process for the preparation of ethyl
4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-
methylamino)pyrimidine-5-carboxylate has been described
25 by Watanabe M. et al., *Bioorg. Med. Chem.* 1997, Vol. 5,
No. 2, 437-444. In this process, in the first stage
p-fluorobenzaldehyde is converted using ethyl iso-
butyrylacetate into an unsaturated ketoester, which is
then cyclocondensed in the second stage with S-
30 methylisothiurea hydrogensulphate and subsequently
dehydrated in the third stage to give the corresponding
pyrimidine. In the fourth stage, this is then oxidized


using m-chloroperbenzoic acid to give the corresponding sulphonylpyrimidine, which is then reacted in the fifth stage with methylamine and subsequent treatment with methanesulphonyl chloride to give ethyl 4-(4-fluoro-phenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methyl-amino)pyrimidine-5-carboxylate.

It is disadvantageous in this process, on the one hand, that many reaction stages are necessary and, on the other hand, that the desired product is only formed in moderate yield.

The object of the invention was to make available an economical, industrially feasible process for the preparation of compounds of the formula I.

The object is achieved by the novel process according to Patent Claim 1.

According to the invention, compounds of the general formula



in which

R is hydrogen or a group of the formula $-\text{SO}_2\text{R}^1$;

R¹ is C₁₋₆-alkyl;


R² is hydrogen or C₁₋₆-alkyl;

R³ is C₁₋₆-alkyl;


R⁴ is C₁₋₆-alkyl,

are prepared in that, in a first stage, a compound of the general formula


- 3 -



in which R³ and R⁴ have the abovementioned meaning, is reacted in the presence of a Lewis acid with 4-fluorobenzonitrile to give a compound of the general
5 formula



in which R³ and R⁴ have the abovementioned meaning, and in a second stage the compound of the formula III obtained is reacted with a compound of the general
10 formula



in which R and R² have the abovementioned meaning, to give the final product of the formula I.

15 "C₁₋₆-alkyl" is understood here and below as meaning linear and branched alkyl groups having 1-6 carbon atoms, such as, for example, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, isobutyl, sec-butyl, pentyl and its isomers and hexyl and its
20 isomers.

The compounds of the formula II can be prepared according to *Chem. Berichte* 1958, 91, 759 or are commercially available organic synthetic chemicals.

- 4 -

4-Fluorobenzonitrile is a commercially available organic synthetic chemical.

5 The Lewis acid employed in the first stage is expediently an aprotic Lewis acid such as, for example, tin tetrachloride, titanium tetrachloride or aluminium chloride. Tin tetrachloride is preferably employed.

10 The first stage is expediently carried out in the presence of an organic solvent. The organic solvents employed can be, for example, aromatic hydrocarbons, chlorinated aromatic and aliphatic hydrocarbons. Aromatic hydrocarbons employed are preferably toluene, benzene or xylene. The chlorinated
15 aromatic hydrocarbon employed is preferably chlorobenzene; the chlorinated aliphatic hydrocarbon employed is preferably 1,2-dichloroethane. Toluene and 1,2-dichloroethane are particularly preferably employed.

20 The reaction in the first stage is expediently carried out at a temperature from -5 to 140°C, advantageously at 60 to 100°C.

25 After a reaction time of 30 min to 6 h and subsequent hydrolysis, the compounds of the formula III can be isolated by known methods such as, for example, by extraction or can be employed directly, without isolation, for the second stage. The intermediate
30 (formula III) is preferably isolated.

Compounds of the formula III include cis and trans isomers.

35 In a second stage, a compound of the formula III is reacted with a compound of the formula IV to give the final product of the formula I.

- 5 -

The invention comprises, on the one hand, compounds of the formula I in which R and R² are hydrogen. These compounds are prepared by reaction of compounds of the formula III with cyanamide.

5

The reaction with cyanamide is expediently carried out in the presence of an organic solvent, a mixture of water with an organic solvent or in water. Water is particularly preferably employed. Organic solvents employed are advantageously toluene or ethyl acetate. Organic solvents employed as a mixture with water are advantageously alcohols such as, for example, methanol, ethers such as, for example, dioxane or aromatic hydrocarbons such as, for example, toluene or *N,N*-dimethylacetamide.


10

The reaction with cyanamide is expediently carried out at a temperature of 10 to 120°C, advantageously at 40 to 100°C. The pH is expediently in a range from 3 to 9, advantageously in a range from 4 to 7. After a reaction time of, in total, 1 to 20h, the compounds of the general formula I are obtained, which can be worked up according to known methods.

20

In a particular embodiment, 2-amino-4-(4-fluorophenyl)-6-isopropylpyrimidine-5-carboxylic acid esters of the general formula


25




in which R³ has the meaning indicated in formula I are prepared in that, in a first stage, an alkyl isobutyrylacetate of the general formula

30


- 6 -



in which R³ has the meaning mentioned is reacted in the presence of a Lewis acid with 4-fluorobenzonitrile to give a 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoic acid ester of the general formula



and in a second stage the compound of the formula IIIa is reacted with cyanamide of the formula




10 in which R and R² are hydrogen, to give the final product of the formula Ia.

The radical R³ is preferably methyl.


15 The compounds of the formula III are novel and also a subject of the invention.

The invention comprises, on the other hand, compounds of the formula I in which R is a group of the formula -SO₂R¹ and R¹ and R² are C₁₋₆-alkyl. These 20 4-(4-fluorophenyl)-6-alkyl-2-(N-alkanesulphonyl-N-alkylamino)pyrimidine-5-carboxylic acid esters of the general formula


- 7 -



in which R¹, R², R³ and R⁴ are identical or different and are a C₁₋₆-alkyl group, can be prepared in that
 5 alkanolic acid esters of the general formula



in which R³ and R⁴ have the abovementioned meaning, are reacted with N-cyano-N-alkylalkanesulphonamides, optionally isolated or prepared in situ, of the general
 10 formula



in which R¹ and R² are a C₁₋₆-alkyl group.

The reaction can be carried out either in the
 15 presence of a base or in the presence of a Lewis acid.

Bases which can be employed are alkali metal compounds such as, for example, alkali metal hydrides, alkali metal carbonates, alkali metal alkoxides or
 20 alkali metal silazanes. Alkali metal carbonates which can be used are lithium, sodium or potassium carbonate. The alkali metal hydride employed can be potassium,

lithium or sodium hydride; sodium hydride is preferably employed. The alkali metal alkoxide employed can be sodium or potassium tert-pentoxide or sodium or potassium tert-butoxide, preferably sodium tert-pentoxide or sodium tert-butoxide. The alkali metal silazane used can be sodium hexamethyldisilazane or potassium hexamethyldisilazane. The base preferably employed is an alkali metal hydride or an alkali metal alkoxide.

10

The reaction is expediently carried out in the presence of a base in a polar organic solvent. The polar solvent used can be, for example, *N,N*-dimethylacetamide, isopropanol, tert-butanol, toluene, dimethylformamide, tetrahydrofuran, 1,4-dioxane or mixtures of these. *N*-Alkylalkanesulphonamides such as, for example, *N*-methylmethanesulphonamide are likewise suitable as solvents. The reaction is preferably carried out in *N*-alkylalkanesulphonamide and tert-butanol.

15
20

The reaction can be carried out in the presence of a base at a temperature from -10 to 150°C, preferably from 0 to 80°C.

25

The reaction of compounds of the formula IIb with compounds of the formula IVb in the presence of a Lewis acid is expediently carried out in the solvent which is inert to the Lewis acid. Inert solvents which can be employed are, for example, aromatic hydrocarbons, and chlorinated aromatic and aliphatic hydrocarbons. Aromatic hydrocarbons employed are preferably toluene or xylene. The chlorinated aromatic hydrocarbon employed is preferably chlorobenzene; chlorinated aliphatic hydrocarbons employed are preferably dichloromethane or 1,2-dichloroethane.

30
35

- 9 -

The reaction in the presence of a Lewis acid can be carried out at a temperature from 20 to 150°C, preferably from 80 to 120°C.

5 Suitable Lewis acids are for example TiCl_4 , TiBr_4 or SnCl_4 . Titanium tetrachloride is preferred.

The amount of Lewis acid is 0.1 to 2 molar equivalents based on the compound of the formula IIIb.

10


After a reaction time of 1 to 24 h, the final products of the general formula Ib can be isolated by known working-up methods.

15

In a preferred embodiment, the final product of the formula Ib isolated is ethyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulphonyl-*N*-methylamino)-pyrimidine-5-carboxylate ($\text{R}^1 = \text{R}^2 = \text{R}^3 = \text{methyl}$, $\text{R}^4 = \text{isopropyl}$).

20

Compounds of the formula IVb are expediently prepared in that a compound of the general formula



25 in which R^1 and R^2 are C_{1-6} -alkyl, is reacted with cyanogen halide in the presence of a base.

Suitable bases are the bases described beforehand.

30

The cyanogen halide employed can be cyanogen fluoride, cyanogen chloride, cyanogen bromide or cyanogen iodide. Cyanogen chloride or cyanogen bromide is preferably employed.

This reaction can likewise be carried out in the polar organic solvents described beforehand. The reaction is preferably carried out in tetrahydrofuran.

5 The reaction is expediently carried out at a temperature from -20 to 50°C, preferably at a temperature from -10 to +20°C.


10 After a reaction time of $\frac{1}{4}$ to 1 h, the compounds of the formula IVb, which have not yet been described in the literature, can then be isolated in a manner known to the person skilled in the art.

15 These compounds of the formula IVb are also a subject of the invention.

20 Compounds of the formula IVb can be prepared in situ, i.e. they are formed directly from the corresponding starting materials during the reaction without isolation. However, they can also be prepared and isolated separately in order then to employ them for the reaction.

25 Examples of compounds of the formula IVb are: N-cyano-N-methylmethanesulphonamide, N-cyano-N-ethylmethanesulphonamide, N-cyano-N-propylmethanesulphonamide, N-cyano-N-butylmethanesulphonamide, N-cyano-N-pentylmethanesulphonamide and N-cyano-N-hexylmethanesulphonamide. N-Cyano-N-methylmethanesulphonamide is
30 preferred.

The compounds of the formula I can also be prepared in that a compound of the general formula



in which R^3 and R^4 have the meaning mentioned in Claim 1, is reacted with a compound of the formula IV.

5 The reaction is carried out analogously to the reaction of the compounds of the formula III with compounds of the formula IV, preferably in the presence of a base in a polar organic solvent at a temperature from -10 to 150°C.

10

Suitable bases and solvents correspond to the bases and solvents which are listed under the reaction of compounds of the formula III with compounds of the formula IV.

15

In a preferred embodiment, compounds of the general formula Ib are prepared in that a compound of the formula VI is reacted with a compound of the formula IVb in a polar organic solvent at a temperature from 0 to 80°C in the presence of a base.

20

The compounds of the formula VI can be prepared by reaction of C_{1-6} -alkyl nitriles with C_{1-6} -alkyl 4-fluorobenzoylacetate in the presence of a Lewis acid.

25 A compound of the formula VI in which R^3 is methyl and R^4 is isopropyl is preferred. The Lewis acid is preferably tin tetrachloride. The reaction is expediently carried out in a polar solvent. Suitable solvents correspond to the solvents which are listed

30 under the reaction of compounds of the formula II with compounds of the formula III described above. The reaction in the first stage is expediently carried out

at a temperature from -5 to 140°C, advantageously at 60 to 100°C.

The compounds of the formula VI are novel and
5 likewise a subject of the invention.

Examples:

Example 1:

10 **Methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate**

IIIa, R³ = methyl, toluene, SnCl₄

1.50 g of methyl isobutyrylacetate (10.0 mmol, concentration 96%) and 1.24 g of 4-fluorobenzonitrile
15 (10 mmol, concentration 98%) were dissolved in 10 ml of toluene and treated with 2.63 g of tin tetrachloride (10 mmol, concentration 99%) at room temperature over the course of 6 min. After 3 h at room temperature, the mixture was heated to 80°C. After 2.5 h, the suspension
20 was again cooled to room temperature and treated with 10 ml of water. It was diluted with 5 ml of ethyl acetate and the phases were separated. After extraction of the aqueous phase with ethyl acetate (2 × 5 ml), the combined organic phases were dried over magnesium
25 sulphate. After concentration in vacuo, 3.50 g of crude product were obtained in the form of a pale yellow, tacky solid. The solid was dissolved in ethyl acetate and purified by flash chromatography (n-hexane/ethyl acetate 5:1 to 1:1). After concentration in vacuo,
30 1.44 g of methyl 2-[1-amino-1-(4-fluorophenyl)-methylene]-4-methyl-3-oxopentanoate were obtained.
Yield: 54.3% (concentration > 99%) in the form of a colourless solid.
Melting point: 85.8 - 86.6°C.

Example 2:**Methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate**IIIa, R³ = methyl, toluene, SnCl₄

5 22.53 g of methyl isobutyrylacetate (0.15 mol, concentration 96%) and 18.54 g of 4-fluorobenzonitrile (0.15 mol, concentration 98%) were dissolved in 150 ml of toluene and treated with 43.32 g of tin tetrachloride (0.165 mol, concentration 99%) at room
10 temperature over the course of 12 min. After half an hour at room temperature, the mixture was heated to 80°C. After 3 h, the suspension was cooled to room temperature and treated with 150 ml of water. It was
15 diluted with 100 ml of ethyl acetate and the phases were separated. After extraction of the aqueous phase with ethyl acetate (2 x 100 ml), the combined organic phases were washed with 100 ml of saturated aqueous sodium hydrogencarbonate solution and 100 ml of 1N
20 sodium hydroxide solution and dried over magnesium sulphate. After concentration in vacuo, 46.46 g of crude product were obtained in the form of a yellowish solid. The solid was dissolved in a mixture of 50 ml of n-hexane and 5 ml of toluene under reflux and filtered hot. The product precipitated from the filtrate on
25 cooling. By filtering through a frit and washing the filter cake with 2 x 40 ml of n-hexane, 32.42 g of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were obtained in the form of a spectroscopically pure solid.

30 Yield: 80.7% (concentration 99.0%)

Melting point: 84.0 - 84.9°C.

¹H NMR (DMSO-d₆, 400 MHz): δ = 0.98 (d, 6H); 3.06 (sept, 1H); 3.24 (s, 3H); 7.27 (t, 2H); 7.35 (m, 2H);
35 8.38 (s, 1H); 10.59 (s, 1H).

¹³C NMR (DMSO-d₆, 100 MHz): δ = 19.45 (q); 36.04 (d); 50.74 (q); 101.54 (s); 115.11 (sd); 115.33 (sd);

129.16 (dd); 129.25 (dd);
133.68 (sd); 161.46.54
(s); 163.91 (s); 165.59
(s); 169.71 (s); 201.10
(s).

5

Example 3**Methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate**

10 IIIa, R³ = methyl, toluene, SnCl₄

75.09 g of methyl isobutyrylacetate (0.50 mol, concentration 96%) and 61.80 g of 4-fluorobenzonitrile (0.50 mol, concentration 98%) were introduced into [500 ml of toluene and treated with 144.7 g of tin tetrachloride (0.55 mol, concentration 99%) at room temperature over the course of 16 min. After half an hour at room temperature, the thick suspension was heated to 80°C. After 3 h, 175 ml of toluene were distilled off at normal pressure, and the mixture was cooled to room temperature and treated with 450 ml of saturated sodium carbonate and 300 ml of ethyl acetate. The organic phase was separated off and the aqueous phase (after dilution with 300 ml of water) was again extracted with 300 ml of ethyl acetate. After drying the combined organic phases over sodium sulphate, the solvent was removed in vacuo (40°C/25 mbar). 127.2 g of crude product were obtained in the form of a slightly yellowish solid, which was dissolved in 160 ml of n-hexane (slightly turbid solution). After hot filtration and subsequent cooling in an ice bath, 99.8 g of methyl 2-[1-amino-1-(4-fluorophenyl)-methylene]-4-methyl-3-oxopentanoate were obtained after filtration in the form of a pale yellow solid.
Yield: 74.5% (concentration 99.3%)
35 Melting point: 86.4 - 87.8°C.

Example 4**Methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate**

IIIa, R^3 = methyl, 1,2-dichloroethane, $AlCl_3$

- 5 754 mg of methyl isobutyrylacetate (5.00 mmol, concentration 96%) and 618 mg of 4-fluorobenzonitrile (5.00 mmol, concentration 98%) were introduced into 5 ml of 1,2-dichloroethane and treated with 673 mg of aluminium chloride (5.00 mol) at room temperature.
- 10 After one hour at room temperature, the mixture was heated to 80°C. After 19 h - the mixture contained 13.9 area per cent of product according to HPLC analysis - the mixture was cooled to room temperature and treated with water (5 ml). The organic phase was
- 15 separated off and the aqueous phase was extracted with dichloromethane (3 x 5 ml). The combined organic phases were dried over magnesium sulphate and concentrated in vacuo. 1.01 g of crude product were obtained, which contained 3% (30 mg) of methyl 2-[1-amino-
- 20 1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate according to 1H -NMR spectrum.

Example 5**Methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate**

IIIa, R^3 = methyl, toluene, $SnCl_4$

- 25 75.09 g of methyl isobutyrylacetate (0.50 mol, concentration 96%) and 61.8 g of 4-fluorobenzonitrile (0.50 mol, concentration 98%) were dissolved in 500 ml
- 30 of toluene and treated with 144.72 g of tin tetrachloride (0.55 mol, concentration 99%) at room temperature over the course of 15 min. After half an hour at room temperature, the mixture was heated to 80°C. After 3 h, the suspension was cooled to 10°C and
- 35 treated with 500 ml of water. The mixture was diluted with 100 ml of dichloroethane and the phases were separated. The organic phase was washed with 2 x 100 ml of 1N sodium hydroxide solution. After concentration in vacuo, 125.5 g of crude product were obtained in the

- 16 -

form of a yellowish solid. The solid was dissolved in 160 ml of n-hexane under reflux and filtered hot. The product precipitated from the filtrate on cooling. By filtering through a frit and washing the filter cake with 130 ml of n-hexane, 79.8 g of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were obtained in the form of a spectroscopically pure solid.

Yield: 60.7% (concentration 98.7%)

10 Melting point: 88.4 - 89.3°C.

Example 6

Methyl 2-amino-4-(4-fluorophenyl)-6-isopropylpyrimidine-5-carboxylate

15 Ia, R³ = methyl, water

A suspension of 1.33 g of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate (5.00 mmol) in 4.20 g of 50% strength aqueous cyanamide solution (50.0 mmol) was heated to reflux. After 17 h, it was cooled to room temperature and treated with ethyl acetate (5 ml) and water (5 ml). The undissolved solid was filtered off. The organic phase was separated off and the aqueous phase was extracted with ethyl acetate (2 x 5 ml). The combined organic phases were dried over magnesium sulphate and concentrated in vacuo. The crude product (0.97 g) was purified by flash chromatography on silica gel (eluent: methylene chloride). 121.6 mg of methyl 2-amino-4-(4-fluorophenyl)-6-isopropylpyrimidine-5-carboxylate were obtained in the form of a colourless solid.

Yield: 8.4%

Melting point: 146.2 - 147.0°C.

¹H NMR (DMSO-d₆, 400 MHz): δ = 1.08 (d, 6H); 3.04 (sept, 1H); 3.57 (s, 3H); 7.05 (s, broad, 2H); 7.28 (m, 2H); 7.53 (m, 2H).

35

Example 7**Methyl 2-amino-4-(4-fluorophenyl)-6-isopropylpyrimidine-5-carboxylate**

Ia, R³ = methyl, *N,N*-dimethylacetamide

- 5 5.00 g of methyl 2-[1-amino-1-(4-fluorophenyl)-methylene]-4-methyl-3-oxopentanoate (18.7 mmol) were treated with 5 ml of *N,N*-dimethylacetamide and 15.71 g of a 50% strength aqueous cyanamide solution (187 mmol) and heated to reflux. After 5 h, the solution was
10 poured onto 50 ml of water. The mixture was cooled in an ice bath and the deposited precipitate was isolated by means of a suction filter. After drying, 2.72 g of crude product were obtained in the form of a yellowish solid. After flash chromatography (150 g of silica gel;
15 eluent: hexane/ethyl acetate 3:2), 1.19 g of methyl 2-amino-4-(4-fluorophenyl)-6-isopropylpyrimidine-5-carboxylate were isolated in the form of a colourless solid.
Yield: 22.0%
20 Melting point: 145 - 146°C.

Example 8***N*-Cyano-*N*-methylemethanesulphonamide**

IVb, R¹ = R² = methyl

- 25 12.22 g (0.28 mol) of sodium hydride (55% in oil) were twice suspended in 100 ml of *n*-hexane under nitrogen and freed from the oil by means of a frit. The hexane-moist sodium hydride was taken up in 200 ml of tetrahydrofuran and cautiously treated at 2°C with
30 22.98 g (0.20 mol) of *N*-methylemethanesulphonamide (concentration 95%). Evolution of gas was observed. After addition was complete (25 min), the cooling bath was removed and the mixture was subsequently reacted at room temperature for 3 h 40 min until evolution of
35 hydrogen was no longer observed. The mixture was subsequently cooled again using an ice bath and 20.0 g (0.32 mol) of cyanogen chloride (concentration 99%) were cautiously introduced over the course of 40 min (slightly exothermic). After subsequent reaction at

- 18 -

0 - 5°C for 45 min, the reaction mixture was poured onto 200 ml of ice water. The phases were separated and the aqueous phase was extracted with 1 x 200 ml and two times 100 ml of diethyl ether. The organic phases were
5 combined and dried over magnesium sulphate. After filtration and concentration in vacuo, 26.57 g of crude product were obtained in the form of a yellowish, two-phase oil, which partially crystallized in the refrigerator. After distillation in vacuo (boiling
10 point 90 - 95°C/0.01 mbar), 17.24 g of *N*-cyano-*N*-methylmethanesulphonamide were obtained in the form of a colourless oil, which crystallized in the refrigerator.

Yield: 61.7%, [concentration: 96.0% (GC)]

15 Melting point: 29 - 30°C.

Example 9

N-Cyano-*N*-methylmethanesulphonamide

IVb, $R^1 = R^2 = \text{methyl}$

20 36.65 g (0.84 mol) of sodium hydride (55% in oil) were suspended three times in 200 ml of *n*-hexane under nitrogen and freed from the oil by means of a frit. The hexane-moist sodium hydride was taken up in 600 ml of tetrahydrofuran and cautiously treated at 2°C
25 with 68.94 g (0.60 mol) of *N*-methylmethanesulphonamide (concentration 95%). Evolution of gas was observed. After addition was complete (45 min), the cooling bath was removed and the mixture was subsequently reacted at room temperature for 4 h 25 min until evolution of
30 hydrogen was no longer observed. It was subsequently cooled again using an ice bath and 59.6 g (0.96 mol) of cyanogen chloride (concentration 99%) were cautiously introduced over the course of 1 h 45 min (slightly exothermic). After subsequent reaction at 0 - 5°C for
35 20 min, the reaction mixture was poured onto 600 ml of ice water. The phases were separated and the aqueous phase was extracted with 2 x 500 ml of diethyl ether. The combined organic phases were dried over magnesium sulphate. After filtration and concentration in vacuo,

99.22 g of crude product were obtained, which crystallized at 15°C. After filtration, 70.56 g of crude product were isolated in the form of a tacky, colourless solid. After distillation in vacuo (boiling point 90 - 95°C/0.01 mbar), 57.4 g of *N*-cyano-*N*-methylmethanesulphonamide were obtained in the form of a colourless oil, which crystallized in the refrigerator. Yield: 71.3%, [concentration: > 99% (GC)]
Melting point: 29.0 - 30.0°C.

¹H NMR (DMSO-d⁶, 400 MHz): δ = 3.28 (s, 3H); 3.48 (s, 3H).
¹³C NMR (DMSO-d⁶, 100 MHz): δ = 35.46 (q); 37.74 (q); 109.35 (s).

15 Example 10

N-Cyano-*N*-methylmethanesulphonamide

IVb, R¹ = R² = methyl

36.7 g (0.84 mol) of sodium hydride (55% in oil) were suspended three times in 200 ml of *n*-hexane under nitrogen and freed from the oil by means of a frit. The hexane-moist sodium hydride was taken up in 600 ml of tetrahydrofuran and cautiously treated at 2°C with 68.94 g (0.60 mol) of *N*-methylmethanesulphonamide (concentration 95%). Evolution of gas was observed. After addition was complete (30 min), the cooling bath was removed and the mixture was subsequently reacted at room temperature for 4 h until evolution of hydrogen was no longer observed. It was subsequently cooled again using an ice bath and 59.6 g (0.96 mol) of cyanogen chloride (concentration 99%) were cautiously introduced over the course of 2 h (slightly exothermic). After subsequent reaction at 0 - 5°C for 20 min, the reaction mixture was poured onto 600 ml of ice water. After addition of 500 ml of diethyl ether, the phases were separated and the aqueous phase was extracted with 500 ml of diethyl ether. The combined organic phases were dried over magnesium sulphate. After filtration and concentration in vacuo, 99.22 g of crude product were obtained, which crystallized at

- 20 -

15°C. After filtration, 86.5 g of crude product were isolated in the form of a tacky, colourless solid. After distillation in vacuo (boiling point 91 - 93°C/0.1 mbar), 72.9 g of *N*-cyano-*N*-methanethanesulphonamide were obtained in the form of a colourless oil which crystallized in the refrigerator. Yield: 90.6%, [concentration: > 99% (GC)]

Example 11

10 **Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulphonyl-*N*-methylamino)pyrimidine-5-carboxylate**
Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, NaH
2.66 g (10.0 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate
15 were introduced into 12 g of *N,N*-dimethylacetamide together with 2.70 g (20.0 mmol) of *N*-cyano-*N*-methanethanesulphonamide. 440 mg (11.0 mmol) of sodium hydride (60% in oil) were added to this solution at room temperature over the course of 2 h. An orange-red
20 clear solution was obtained. After 6 h at room temperature, the reaction solution was poured onto 25 ml of water. The suspension was stirred in an ice bath for 30 minutes, and the precipitate was filtered off and washed with water (2 x 10 ml). After drying in
25 a high vacuum, 1.11 g (29.1%) of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulphonyl-*N*-methylamino)pyrimidine-5-carboxylate were obtained in the form of a pale beige solid. The product still contained traces of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate.
30 ^1H NMR (DMSO- d_6 , 400 MHz): $\delta = 1.15$ (d, 6H); 3.17 (sept, 1H); 3.50 (s, 3H); 3.58 (s, 3H); 3.73 (s, 3H); 7.39 (m, 2H); 7.69 (m, 2H).
35

Example 12

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, sodium tert-pentoxide

2.66 g (10.0 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 7 g of *N,N*-dimethylacetamide together with 2.70 g (20.0 mmol) of *N*-cyano-*N*-methylmethanesulphonamide. 1.21 g (11.0 mmol) of sodium tert-pentoxide in 5 g of dimethylacetamide were added to this solution at room temperature after the course of 3 h. An orange-red clear solution was obtained. After 2.5 h at room temperature, the reaction solution was poured onto 25 ml of water. The suspension was stirred in an ice bath for 30 minutes, and the precipitate was filtered off and washed with water (10 ml). After drying in a high vacuum, 1.65 g of crude product were obtained in the form of a pale beige solid which contained 676.5 mg of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulphonyl-*N*-methylamino)-pyrimidine-5-carboxylate.
Yield: 17.7%

Example 13

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, NaH

10.0 g (37.7 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 45 ml of *N,N*-dimethylacetamide together with 15.2 g (113 mmol) of *N*-cyano-*N*-methylmethanesulphonamide. 3.50 g (88.7 mmol) of sodium hydride (60% in oil) were added at room temperature to this solution. An orange-red viscous solution was obtained. The reaction solution was poured onto 120 ml of water and stirred in an ice bath for 30 minutes. The precipitate was filtered off and washed with water (10 ml). After drying in a high vacuum, 4.60 g of crude

- 22 -

product were obtained in the form of a pale beige solid. According to the ^1H NMR spectrum, the solid contained an 84 : 16 product/starting material mixture, which corresponds to a yield of 27% of methyl
5 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulphonyl-*N*-methylamino)pyrimidine-5-carboxylate.

Example 14

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methane-
10 sulphonyl-*N*-methylamino)pyrimidine-5-carboxylate

Ib, $\text{R}^1 = \text{R}^2 = \text{R}^3 = \text{methyl}$, $\text{R}^4 = \text{isopropyl}$, NaH, compound of the formula IVb formed in situ

2.65 g (10.0 mmol) of methyl 2-[1-amino-
1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate
15 were introduced into 5 g of *N,N*-dimethylacetamide together with 2.30 g (20.0 mmol) of *N*-methylmethane-sulphonamide. 470 mg (20.0 mmol) of sodium hydride (60% in oil) were added to this solution at room temperature. The reaction solution was treated with
20 2.40 g (40.0 mmol) of cyanogen chloride at room temperature. After 20 h, the mixture was poured onto 40 ml of water and the yellowish suspension was cooled in an ice bath. The precipitate deposited was filtered off and washed with 20 ml of water. After drying in
25 vacuo, 2.22 g of crude product were obtained in the form of a pale beige solid, which according to the ^1H NMR spectrum contained a 70 : 30 starting material/product mixture, which corresponds to a yield of 22% of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulphonyl-*N*-methylamino)pyrimidine-5-
30 carboxylate.

Example 15

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methane-
35 sulphonyl-*N*-methylamino)pyrimidine-5-carboxylate

Ib, $\text{R}^1 = \text{R}^2 = \text{R}^3 = \text{methyl}$, $\text{R}^4 = \text{isopropyl}$, NaH, compound of the formula IVb formed in situ

13.3 g (50.0 mmol) of methyl 2-[1-amino-
1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate

- 23 -

were introduced into 25 g of *N,N*-dimethylacetamide together with 10.9 g (100.0 mmol) of *N*-methylethanesulphonamide. 3.60 g (90.0 mmol) of sodium hydride (60% in oil) were added to this solution at 5 25°C. Vigorous foaming was observed. After 15 min at 25°C, 6.0 g (100.0 mmol) of cyanogen chloride gas were introduced after the course of 20 min. An orange-coloured suspension was formed. This was stirred at 25°C for 2 h and 2.40 g (60.0 mmol) of sodium hydride 10 (60% in oil) and subsequently 6.0 g (100.0 mmol) of cyanogen chloride were added again. The mixture was stirred at 25°C again for 1 h before a further 3.60 g (90.0 mmol) of sodium hydride (60% in oil) were added. The reaction mixture was poured onto an ice/water 15 mixture (200 ml) and stirred at 0°C for 1 h. The resulting solid was filtered off and washed with 100 ml of water. After drying in a high vacuum, 14.62 g of crude product were obtained in the form of a beige solid. 8.22 g of this solid were recrystallized in an 20 acetone/water mixture. 4.24 g of product were obtained in the form of a pale beige solid, which corresponds to a yield of 29% [concentration (GC) 72%].

Example 16

25 **Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulphonyl-*N*-methylamino)pyrimidine-5-carboxylate**

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, NaH, compound of the formula IVb formed in situ

13.3 g (50.0 mmol) of methyl 2-[1-amino-30 1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 25 g of *N,N*-dimethylacetamide together with 10.9 g (100.0 mmol) of *N*-methylethanesulphonamide. 3.60 g (90.0 mmol) of sodium hydride (60% in oil) were added to this solution at 35 25°C. Vigorous foaming was observed. After 25 min at 25°C, 6.0 g (100.0 mmol) of cyanogen chloride gas were introduced after the course of 12 min. During this, the temperature rose temporarily to 32°C. An orange-coloured suspension was formed. The mixture was stirred

- 24 -

at 25°C for 1 h 25 min and 2.40 g (60.0 mmol) of sodium hydride (60% in oil) and subsequently 3.0 g (50.0 mmol) of cyanogen chloride were added again. The mixture was stirred at 25°C again for 1 h 40 min before a further
5 2.40 g (60.0 mmol) of sodium hydride (60% in oil) and 3.0 g (50.0 mmol) of cyanogen chloride were added. After addition of a further 2.40 g (60.0 mmol) of sodium hydride (60% in oil), the reaction mixture was poured onto an ice/water mixture (200 ml). The
10 apparatus was washed with 50 ml of water and the mixture was stirred at 0°C for 1 h. The resulting solid was filtered off and washed with 50 ml of water. After drying in a high vacuum, 10.47 g of crude product were obtained in the form of a beige solid. 8.00 g of this
15 solid were recrystallized in an acetone/water mixture. 5.50 g of product were obtained in the form of a pale beige solid, which corresponded to a yield of 30.2% [concentration (GC) 80%].

20 Example 17

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, NaH, compound of the formula IVb formed in situ

25 13.3 g (50.0 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 50.0 g (0.46 mol) of N-methyl-methanesulphonamide. 12.0 g (0.30 mol) of sodium hydride (60% in oil) were added to this solution at
30 25°C after the course of 1 h 45 min. Vigorous foaming was observed. In order to guarantee better stirrability, 30.0 g (0.275 mol) of N-methyl-methanesulphonamide were added again during the addition. After 30 min at 25°C, 15.0 g (0.25 mol) of cyanogen
35 chloride gas were introduced over the course of 50 min. The suspension was transferred to an autoclave and stirred at 60°C for 18.5 h. The reaction mixture was poured onto an ice/water mixture (200 ml) and stirred at 0°C for 30 min. The resulting solid was filtered off

- 25 -

and washed with 50 ml of water. After drying in a high vacuum, 18.71 g of crude product were obtained in the form of a beige solid. 10.0 g of this solid were recrystallized in an acetone/water mixture. 5.40 g of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate were obtained in the form of a colourless solid. Yield of 50.6%, [concentration 95.5% (GC)].

10 Example 18

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, sodium tert-pentoxide, compound of the formula IVb formed in situ

15 13.3 g (50.0 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 50.0 g (0.46 mol) of N-methylmethanesulphonamide. 33.0 g (0.30 mol) of sodium tert-pentoxide were added to this solution at 25°C

20 after the course of 20 min. A yellowish, thick suspension was formed. After 30 min at 25°C, 15.0 g (0.25 mmol) of cyanogen chloride gas were introduced after the course of 25 min. The suspension, which was now more readily stirrable, was transferred to an

25 autoclave and stirred at 60°C for 17 h. The reaction mixture was poured onto an ice/water mixture (200 ml) and stirred at 0°C for 30 min. The resulting solid was filtered off and washed with 50 ml of water. After drying in a high vacuum, 20.81 g of crude product were

30 obtained in the form of a beige solid having a concentration of about 68% (GC). This corresponded to a yield of 74.1% of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate.

Example 19

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, TiCl_4 ,
5 chlorobenzene

10.0 g (37.7 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 50 ml of chlorobenzene together with 10.17 g (75.4 mmol) of N-cyano-N-methylmethanesulphonamide and the mixture was treated at room temperature with 7.22 g (37.7 mmol) of titanium tetrachloride. The reaction progressed exothermically. The red-orange-coloured suspension was heated to 110 - 120°C and stirred for 3.5 h. It was then cooled to room temperature and treated with 30 ml of water. The organic phase was separated off and the aqueous phase was extracted with methylene chloride (30 ml). The combined organic phases were washed with 30 ml of water and dried over magnesium sulphate. After filtering and concentrating the solution in a water-jet vacuum and drying in vacuo, 11.48 g of crude product were obtained in the form of a tacky solid.
Yield: 37.8% [concentration (HPLC) 47.3%].

Example 20

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, TiCl_4 ,
chlorobenzene

10.0 g (37.7 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 50 ml of chlorobenzene together with 10.17 g (75.4 mmol) of N-cyano-N-methylmethanesulphonamide and the mixture was treated at room temperature with 14.45 g (75.4 mmol) of titanium tetrachloride. The reaction progressed exothermically. The red-orange-coloured suspension was heated to 110°C and stirred for 17.5 h. It was then cooled to room temperature and treated with 30 ml of water. The

organic phase was separated off and the aqueous phase was extracted with methylene chloride (2 x 30 ml). The combined organic phases were washed with 30 ml of water and dried over magnesium sulphate. After filtering and
5 concentrating the solution in a water-jet vacuum and drying in vacuo, 12.92 g of crude product were obtained in the form of a brownish oil.
Yield: 23.3% [concentration (HPLC) 25.9%].

10 Example 21

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate
Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, TiCl_4 , toluene
10.0 g (37.7 mmol) of methyl 2-[1-amino-
15 1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 50 ml of toluene together with 10.17 g (75.4 mmol) of N-cyano-N-methylmethanesulphonamide and the mixture was treated at room temperature with 3.61 g (18.9 mmol) of titanium tetrachloride. The
20 reaction progressed exothermically. The red-orange-coloured suspension was heated to 110°C and stirred for 4.5 h. It was then cooled to room temperature and treated with 30 ml of water. The organic phase was separated off and the aqueous phase was extracted with
25 ethyl acetate (30 ml). The combined organic phases were washed with 30 ml of water and dried over magnesium sulphate. After filtering and concentrating the solution in a water-jet vacuum and drying in vacuo, 11.54 g of crude product were obtained in the form of a
30 tacky oil.
Yield: 13.5% [concentration (HPLC) 16.8%].

Example 22

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate
35 Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, TiCl_4 , chlorobenzene
10.0 g (37.7 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate

- 28 -

were introduced into 50 ml of chlorobenzene together with 5.08 g (37.7 mmol) of *N*-cyano-*N*-methylethanesulphonamide and the mixture was treated at room temperature with 7.22 g (37.7 mmol) of titanium tetrachloride. The reaction progressed exothermically. The red-orange-coloured suspension was heated to 110 - 120°C and stirred for 5 h. It was then cooled to room temperature and treated with 30 ml of water. The organic phase was separated off and the aqueous phase was extracted with methylene chloride (2 x 30 ml). The combined organic phases were washed with 30 ml of water and dried over magnesium sulphate. After filtering and concentrating the solution in a water-jet vacuum and drying in vacuo, 12.02 g of crude product were obtained in the form of a tacky oil. Yield: 15.0% [concentration (HPLC) 17.9%].

Example 23

Methyl 3-amino-2-[1-(4-fluorophenyl)methanoyl]-4-methylpent-2-enoate

VI, $R^3 = R^4 = \text{methyl}$

5.00 g of methyl 4-fluorobenzoylacetate (24.2 mmol, concentration 95%) and 1.69 g of isobutyronitrile (24.2 mmol, concentration 98%) were dissolved in 25 ml of toluene and treated with 7.01 g of tin tetrachloride (26.6 mmol, concentration >99%) at room temperature after the course of 10 minutes. After 3 h at room temperature, the mixture was heated to 80°C. After 11.5 h, the suspension was again cooled to room temperature and treated with 25 ml of water. The mixture was diluted with 10 ml of ethyl acetate and the phases were separated. The organic phase was washed twice with 10 ml of 1N sodium hydroxide solution and concentrated in vacuo (40°C/25 mbar) after drying over magnesium sulphate. 6.18 g of crude product were obtained in the form of a yellow oil. After chromatography on silica gel (eluent: hexane/ethyl acetate 1.5:1), 3.84 g of methyl 3-amino-2-[1-

- 29 -

(4-fluorophenyl)methanoyl]-4-methylpent-2-enoate were obtained in the form of an oil.

Yield: 59.0%; concentration (HPLC): 98.6%.

GC-MS: $M^+ = 265$.

- 5 ^1H NMR (CDCl_3 , 400 MHz): $\delta =$ 0.98 (d); 1.25 (d); 2.90 (sept); 3.34 (s); 3.47 (s); 3.52 (sept); 5.50 (s, broad); 5.90 (s, broad); 7.08 (m); 7.50 (dd); 7.85 (dd); 9.04 (s, broad); 10.86 (s, broad).
- 10 ^{13}C NMR (CDCl_3 , 100 MHz): $\delta =$ 20.68 (q); 20.76 (q); 30.58 (d); 30.78 (d); 50.57 (q); 51.17 (q); 96.86 (s); 101.09 (s); 114.85 (d); 115.06 (d); 115.16 (d); 115.37 (d); 128.67 (d); 128.76 (d); 131.33 (d); 131.42 (d); 136.98 (s); 137.01 (s); 139.37 (s); 139.40 (s); 162.48 (s); 164.04 (s); 164.95 (s); 166.56 (s); 169.33 (s); 170.33 (s); 171.58 (s); 175.91 (s); 193.24 (s); 194.40 (s).
- 15
- 20
- 25

Example 24

- 30 Methyl 3-amino-2-[1-(4-fluorophenyl)methanoyl]-4-methylpent-2-enoate

VI, $\text{R}^3 = \text{R}^4 = \text{methyl}$

- 35 39.2 g of methyl 4-fluorobenzoylacetate (0.20 mmol) and 16.8 g of isobutyronitrile (0.24 mol, concentration >99%) were dissolved in 200 ml of toluene and treated with 57.9 g of tin tetrachloride (0.22 mmol, concentration >99%) at room temperature after the course of 10 minutes. After 30 min at room temperature, the mixture was heated to 80°C. After 8 h,

- 30 -

the suspension was again cooled to room temperature and treated with 200 ml of water. The mixture was diluted with 200 ml of ethyl acetate and the phases were separated. The organic phase was washed twice with 40 ml of 1N sodium hydroxide solution and concentrated in vacuo (40°C/25 mbar) after drying over magnesium sulphate. 50.9 g of crude product were obtained in the form of a yellow oil. After chromatography of 12.0 g of crude product on silica gel (eluent: hexane/isopropanol 90:10), 10.08 g of methyl 3-amino-2-[1-(4-fluorophenyl)methanoyl]-4-methylpent-2-enoate were obtained in the form of a yellowish oil.
Yield: 86.3%; (concentration: >99%).

Example 25

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate
Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$
2.65 g (10.0 mmol) of methyl 3-amino-2-[1-(4-fluorophenyl)methanoyl]-4-methylpent-2-enoate were introduced into 5.52 g (50.0 mmol) of N-methylmethanesulphonamide and 1.48 g (20.0 mmol) of tert-butanol. 4.95 g (50.0 mmol) of sodium tert-butoxide were added to this solution after the course of 2 min between 23 and 52°C. A yellowish, thick suspension was formed. After 1 h, 2.00 g (32.5 mmol) of cyanogen chloride gas were introduced at 25°C after the course of 20 min. The suspension, which was now more readily stirrable, was stirred for 18 h at 60°C. The reaction mixture was poured onto water (20 ml). The resulting solid was filtered off and washed with water (2 x 5 ml). After drying in a high vacuum, 740 mg of crude product were obtained in the form of a beige solid having a concentration of 70.0% (HPLC). This corresponded to a yield of 13.6% of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate.

Example 26

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$

- 5 2.50 g (9.42 mmol) of methyl 3-amino-2-[1-(4-fluorophenyl)methanoyl]-4-methylpent-2-enoate were introduced into 2.08 g (28.3 mmol) of N-methylmethanesulphonamide and 3.49 g (47.1 mmol) of tert-butanol together with 3.79 g (28.3 mmol) of N-cyano-N-methylmethanesulphonamide. 1.87 g (18.8 mmol) of sodium tert-butoxide were added to this suspension at room temperature after the course of 15 min (exothermic). After 4 h at 76°C, the reaction suspension was poured onto 20 g of ice water. The suspension was stirred in an ice bath, and the precipitate⁷ was filtered off and washed with water (2 x 2.5 ml). After drying in a high vacuum, 1.61 g of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate were obtained in the form of a pale beige solid. Yield: 9.8%; concentration (HPLC): 21.9%.

Example 27

Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

- 25 Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, sodium tert-butoxide, compound of the formula IVb formed in situ 132.6 g (0.50 mol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 276.2 g (2.50 mol) of N-methylmethanesulphonamide and 74.1 g (1.00 mol) of tert-butanol. 243.7 g (2.50 mol) of sodium tert-butoxide were added to this suspension at 30°C in portions such that the temperature did not rise above 60°C. A yellowish, thick suspension was formed. After 20 min at 28°C, the mixture was heated to 50°C and 100.0 g (1.63 mol) of cyanogen chloride gas were introduced after the course of 1 h. The suspension, which was now more readily stirrable, was stirred at

- 32 -

60°C for 19.5 h. The reaction mixture was poured onto water (750 ml) and stirred at room temperature for 15 min. The resulting solid was filtered off and washed with 2 x 250 ml of water and 2 x 200 ml of cold methanol. After drying in a high vacuum, 119.1 g of crude product were obtained in the form of a beige solid having a concentration of 90.1% (HPLC). This corresponded to a yield of 56.3% of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate.

Example 28


Methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate

Ib, $R^1 = R^2 = R^3 = \text{methyl}$, $R^4 = \text{isopropyl}$, NaOtBu, MMSA, CMMSA

2.97 g (11.2 mmol) of methyl 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoate were introduced into 2.45 g (22.4 mmol) of N-methylmethanesulphonamide (MMSA) and 4.19 g (55.9 mmol) of tert-butanol together with 4.51 g (36.6 mmol) of N-cyano-N-methylmethanesulphonamide (CMMSA). 2.22 g (22.4 mmol) of sodium tert-butoxide were added to this suspension at room temperature in portions. An orange-coloured suspension was obtained. It was heated to 50°C and stirred at 50°C for 19.9 h. The suspension was poured onto 20 g of ice water, and the precipitate was filtered off and washed with water (2 x 5 ml). After drying in a high vacuum, 2.91 g of methyl 4-(4-fluorophenyl)-6-isopropyl-2-(N-methanesulphonyl-N-methylamino)pyrimidine-5-carboxylate were obtained in the form of a pale beige solid having a concentration of 76.4% (HPLC), which corresponds to a yield of 52.1%.

Patent Claims:

1. Process for the preparation of compounds of the general formula



5

in which

R is hydrogen or a group of the formula $-\text{SO}_2\text{R}^1$;


R^1 is C_{1-6} -alkyl;

R^2 is hydrogen or C_{1-6} -alkyl;


10 R^3 is C_{1-6} -alkyl;

R^4 is C_{1-6} -alkyl,

characterized in that, in a first stage, a compound of the general formula




- 15 in which R^3 and R^4 have the abovementioned meaning, is reacted in the presence of a Lewis acid with 4-fluorobenzonitrile to give a compound of the general formula



- 20 in which R^3 and R^4 have the abovementioned meaning, and in a second stage the compound of the formula III


- 34 -

obtained is reacted with a compound of the general formula




in which R and R² have the abovementioned meaning, to
5 give the final product of the formula I.

2. Process for the preparation of 2-amino-4-(4-fluorophenyl)-6-isopropylpyrimidine-5-carboxylic acid esters of the general formula




10

in which R³ has the meaning indicated in Claim 1, characterized in that, in a first stage, an alkyl isobutyrylacetate of the general formula




15 in which R³ has the meaning indicated in Claim 1, is reacted in the presence of a Lewis acid with 4-fluorobenzonitrile to give a 2-[1-amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoic acid ester of the general formula

- 35 -



in which R³ has the meaning mentioned, and in a second stage the compound of the formula IIIa is reacted with cyanamide of the formula




- 5 in which R and R² are hydrogen, to give the final product of the formula Ia.
3. Process according to Claim 2, characterized in
10 that R³ is a methyl group.
4. Process according to Claim 2 or 3,
characterized in that the Lewis acid employed in the
first stage is tin tetrachloride.
- 15 5. Process according to one of Claims 2 to 4,
characterized in that the first stage is carried out in
the presence of an organic solvent.
- 20 6. Process according to one of Claims 2 to 5,
characterized in that the reaction in the first stage
is carried out at a temperature from -5 to 140°C.
- 25 7. Process according to one of Claims 2 to 6,
characterized in that the second stage is carried out
in the presence of an organic solvent, a mixture of
water with an organic solvent or in water.

- 36 -

8. Process according to one of Claims 2 to 7, characterized in that the reaction in the second stage is carried out at a temperature from 10 to 120°C.

5 9. Process according to one of Claims 2 to 8, characterized in that the intermediate of the formula IIIa is isolated.


10 10. 2-[1-Amino-1-(4-fluorophenyl)methylene]-4-methyl-3-oxopentanoic acid esters of the general formula



in which R³ has the meaning mentioned in Claim 1.


15 11. Methyl 2-[1-amino-1-(4-fluorophenyl)-methylene]-4-methyl-3-oxopentanoate.

12. Process for the preparation of 4-(4-fluorophenyl)-6-alkyl-2-N-alkansulphonyl-N-alkylamino)-
20 pyrimidine-5-carboxylic acid esters of the general formula




in which R¹, R², R³ and R⁴ are identical or different and are a C₁₋₆-alkyl group, characterized in that a 2-[1-amino-1-(4-fluorophenyl)methylene]-4-alkyl-3-oxo-
25 alkanolic acid ester of the general formula

- 37 -



in which R^3 and R^4 are a C_{1-6} -alkyl group, is reacted with an N-cyano-N-alkylalkanesulphonamide, optionally isolated or prepared in situ, of the general formula



5 in which R^1 and R^2 are a C_{1-6} -alkyl group, to give the final product of the formula Ib.

10 13. Process according to Claim 12, characterized in that the reaction is carried out in a polar organic solvent in the presence of a base.

15 14. Process according to Claim 12 or 13, characterized in that the reaction is carried out at a temperature from -10 to 150°C .


15. Process according to Claim 12, characterized in that the reaction is carried out in an inert solvent in the presence of a Lewis acid.

20 16. Process according to Claim 15, characterized in that the reaction is carried out in the presence of titanium tetrachloride.


25 17. Process according to Claim 15 or 16, characterized in that the reaction is carried out at a temperature from 20 to 150°C .

30 18. Process for the preparation of N-cyano-N-alkylalkanesulphonamides of the general formula

- 38 -




in which R¹ and R² are a C₁₋₆-alkyl group characterized in that a cyanogen halide is prepared using an N-alkyl-alkanesulphonamide of the general formula



5 in which R¹ and R² are a C₁₋₆-alkyl group, in the presence of a base.


19. N-Cyano-N-alkylalkanesulphonamides of the
10 general formula



in which R¹ and R² are a C₁₋₆-alkyl group.

20. N-Cyano-N-methylmethanesulphonamide.

15 21. Process for the preparation of compounds of the general formula I, where R, R¹, R², R³ and R⁴ have the meaning indicated in Claim 1, characterized in that a compound of the general formula



in which R^3 and R^4 have the meaning mentioned in Claim 1, is reacted with a compound of the formula IV.

22. Process for the preparation of a compound of the general formula Ib, characterized in that a compound of the formula VI according to Claim 21 is reacted in the presence of a base with a compound of the formula IVb in a polar organic solvent at a temperature from -5 to 140°C.

23. Compounds of the formula VI, in which R^3 and R^4 have the meaning mentioned in Claim 1.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/06099

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/42 C07C229/34 C07C311/05

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WATANABE M ET AL: "SYNTHESIS AND BIOLOGICAL ACTIVITY OF METHANESULFONAMIDE PYRIMIDINE-AND N-METHANESULFONYL PYRROLE-SUBSTITUTED 3,5-DIHYDROXY-6-HEPTENOATES, A NOVEL SERIES OF HMG-COA REDUCTASE INHIBITORS" BIOORGANIC & MEDICINAL CHEMISTRY, GB, ELSEVIER SCIENCE LTD, vol. 5, no. 2, 1997, pages 437-444, XP000882043 ISSN: 0968-0896 cited in the application page 439, scheme 1	1, 2, 10, 11
A	EP 0 521 471 A (SHIONOGI & CO) 7 January 1993 (1993-01-07) Reference Examples 1, 4	1, 2, 10, 11
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

16 November 2000

Date of mailing of the international search report

21/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

Internation Application No
PCT/EP 90/06099

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 018, no. 656 (C-1286), 13 December 1994 (1994-12-13) & JP 06 256318 A (SHIONOGI & CO LTD), 13 September 1994 (1994-09-13) cited in the application abstract	1,2,21
A	BREAUX E J ET AL: "An improved general synthesis of 4-aryl-5-pyrimidinecarboxylates" JOURNAL OF HETEROCYCLIC CHEMISTRY,US,HETEROCORPORATION. PROVO, vol. 18, January 1981 (1981-01), pages 183-184, XP002117468 ISSN: 0022-152X page 183, scheme I and table I	21,22
A	US 4 013 706 A (ANATOL JESUS ET AL) 22 March 1977 (1977-03-22) columns 7, 8, examples 16, 17	19,20
A	DE 37 17 815 A (BAYER AG) 15 December 1988 (1988-12-15) claim 1	19,20
A	KIN-YA AKIBA ET AL.: "Ring Transformation Equilibrium (Bond Switch) in the 5-(2-Aminovinyl)isothiazole System via Hypervalent Sulfurane. Synthesis, Structure Determination, and Kinetic Study" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 107, no. 9, 1985, pages 2721-30, XP002153123 page 2728, left-hand column, lines 31 and 32 of text (compound 8)	10,11

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Classification No

PCT/EP 88/06099

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0521471	A	07-01-1993	CA 2072945 A	02-01-1993
			HU 61531 A	28-01-1993
			JP 2648897 B	03-09-1997
			JP 5178841 A	20-07-1993
			KR 9605951 B	06-05-1996
			US 5260440 A	09-11-1993
JP 06256318	A	13-09-1994	NONE	
US 4013706	A	22-03-1977	FR 2271202 A	12-12-1975
			FR 2288735 A	21-05-1976
			CH 605700 A	13-10-1978
			DE 2454107 A	22-05-1975
			DE 2462011 A	26-06-1975
			GB 1476796 A	16-06-1977
			GB 1476797 A	16-06-1977
			JP 1197247 C	21-03-1984
			JP 50083331 A	05-07-1975
			JP 58031341 B	05-07-1983
DE 3717815	A	15-12-1988	JP 63307851 A	15-12-1988